

Flexible Sigmoidoscopy

Flex Sig With Magnesium Citrate- Procedure Instructions

Procedure Date: _____ Check-in Time: _____ GI Physician: _____

Location: _____ Procedure Time: _____

Special Instructions:

To ensure a successful exam, please follow all instructions on this sheet carefully. Failure to do so could result in your procedure being postponed or cancelled.

If you are pregnant, please discuss the risks and benefits of this procedure with your physician.

If you take anti-coagulants / blood thinners (for example: Coumadin, Plavix, Pradaxa, Xarelto, Eliquis, Effient, Brilinta, etc.) or Insulin, please notify our office at least 10 days prior to your procedure.

Insulin and other diabetic medications should be reviewed with the prescribing doctor and taken according to their instructions.

Please inform us if you weigh greater than 250 lbs., have kidney disease, congestive heart failure, artificial heart valves, a pacemaker or defibrillator (AICD), sleep apnea, or ascites.

Transportation:

- Please make arrangements to have someone drive you to and from the designated location for your procedure.
- You will not be allowed to use any form of public transportation following your procedure, which includes: Taxi, Bus, Ride-share service, Uber, Lyft and the like.
- The test may be cancelled if you do not have a driver, there will be no exceptions.
- You will be under sedation during your procedure and will be unable to work, drive or operate heavy machinery the day of your procedure.

Flexible Sigmoidoscopy With Magnesium Citrate:

You may obtain Fleet enema and Magnesium Citrate at any pharmacy. IT IS VERY IMPORTANT THAT YOU REFER TO THESE PREP INSTRUCTIONS FROM YOUR DOCTOR REGARDING WHEN YOU NEED TO STOP EATING AND DRINKING. DO NOT FOLLOW THE INSTRUCTIONS THAT YOU RECEIVE FROM THE PHARMACY OR WHAT IS ON THE PREP PACKAGE.

- If you have any questions call our patient line at **1-855-455-4278**
- All heart and blood pressure medications should be continued, including the day of your procedure with small sip of water.
- Insulin and other diabetic medications should be reviewed with the prescribing doctor and taken according to their instructions.

3-Days Before Your Procedure:

- Buy an over-the-counter double Fleet Enema package or two single Fleet Enemas.
- Buy One 300-mL (about 10oz) bottle of Magnesium Citrate.

Day Before Your Procedure:

- On the evening before the procedure, drink the bottle of Magnesium Citrate. Drink two 8oz glasses of water over the next hour.
- Do NOT use any tobacco products or marijuana, do NOT take non-prescribed drugs, and do NOT drink alcohol after midnight before your procedure.

Morning Of Your Procedure:

- Do not eat any food or drink (except a small amount of water for any required morning medications).
- Administer 2 Fleet enemas 30-60 minutes apart, with the second enema being one hour PRIOR TO LEAVING for the examination (to allow appropriate evacuation time).
- In the morning, you should use inhalers, and take heart, blood pressure, seizure, anti-anxiety, and respiratory medications with a sip of water, enough to swallow them (no blood thinners or oral diabetic medications).
- You should wake up early and take these at least 2 hours before your test if possible.

IF sedation is being used:

- Arrange to have someone with you at the end of the exam that can drive you home. A taxi driver alone is NOT sufficient. You will be discharged approximately 60 minutes after your procedure is started.
- Female patients of childbearing age will be required to provide a urine specimen for a pregnancy test prior to anesthesia the day of the procedure. Please avoid urinating prior to arrival the day of your procedure to prevent delay or possible cancellation.